

UDK:582.282.23:618.15 -002-828-02-08

CAUSING FACTORS, SYMPTOMS, TREATMENT AND PREVENTION OF VAGINAL CANDIDIASIS IN MEDICINE

Nabiyeva M.I.

Student of the 1st Medical Faculty of TTA,

dr13liza@gmail.com

Yodgorova N.T.

Associate Professor of the Department of Microbiology, Virology, Immunology, TTA,

tel.yodgorovanodira8@gmail.com, ORCID ID: 0009-0002-4125-4317

Abstract:In the analysis of this literature, general information about vaginal candidiasis, the relevance of the disease, symptoms, treatment and prevention, the fungus that causes this disease types, in particular, general information about the type of *Candida albicans*, etiology, diagnosis of this type about, How does acquired immunodeficiency virus (HIV) cause infectious vaginosis stimulation, the importance of the reduction of probiotics in pregnant women in the development of the disease, the estrogen hormone in women effect of the frequency of change on the disease, different in the metabolism of acids the effect of changes on *Candida albicans*, resistance of the species to various substances, vaccination work, this fungus and o on the disease conducted experiences of recent years, in particular, the effect of antifungal drugs such as nystatin, amphotericin, fluconazole, clotrimazole, their reception and according to instructions and so in theoretical and practical medicine in this field news of recent years information about planned and analyzed and the results are widely reported.

The keywords: Vaginalcandidiasis, *Candida albicans*, HIV, probiotics, antifungal, fluconazole, clotrimazole, infection

Introduction:Among the vaginal complaints in women of reproductive age, yeast infection is the main part of the common diseases vulvovaginal candidiasis caused by a certain type of it.

Incidence and epidemiology:In the last 20 years, the incidence of vulvovaginal candidiasis (VVK) is clear tend to be. Currently, VVK is the second o after bacterial vaginosis stands in line(1). According to different researchers 15 to 40 percent of infectious diseases of the vulva and vagina are caused by a fungal infection. About 75 percent of women Part p is at least once in a lifetime also suffers from this disease. 40-45% of patients have two (or more) VVKs during their lifetime episodes occur10-20% of women are yeast are asymptomatic carrier, fungi often localized in the vagina; fungal colonization can be as high as 40% in pregnant women(34).Epidemiological investigations show that showed that the prevalence of chronic vulvovaginal candidiasis can occur in 7-8% of women who have experienced the first episode. Connect with VVK severe discomfort; significantly reduces the quality of life of young women. Cause candida widespread of vaginal infections and the development of resistance to existing drugs require the need for new special antifungal agents and new targeted drugs. In addition, genetic predisposition in individual patients is also taken into account(8).

The most common symptoms of the disease include vaginal itching, pain during urination, thick white discharge that usually does not smell bad, and redness around the vagina (20). In addition to the above symptoms, vaginal erythema (redness), vaginal fissures (skin cracks), edema (fluid, swelling), as well as in some cases, ulcers are formed around it includes (11) . *Candida albicans* is associated with vaginal candidiasis the most common fungus that type although it is a different type of fungus are the same can cause similar symptoms. Confirmed vaginal candidiasis infections 370 patients in Hungary The study identified the following types of infection:

-Bag to *Candida albicans* candidiasis: 85.7%

- *Candida albicans* (8 types): 13.2% (*C. glabrata*, *C. krusei*, *C. kefir*, *C. parapsilosis*, *C. tropicalis*, *C. dubliniensis*, *C. guilliermondii*, *C. orthopsilosis*) (19).

So, in more than 85% of cases, vulvovaginal candidiasis is primarily caused by *C. Albicans* called This species is a conditional pathogen for the organism. It can cause pathogenicity in the following cases

- Taking drugs (antibiotics, hormonal drugs, cytostatics, oral contraceptives)
- Imbalance of female sex hormones (pregnancy, menopause)
- Type 1 or 2 diabetes, obesity, impaired carbohydrate metabolism, thyroid dysfunction
- Immunosuppression (cancer, HIV infection)
- Avitaminosis that lowers the body's immunity
- Physical exertion and hypothermia
- Stress (33).

Symptoms often increase when women have periods, before puberty (29). Vaginal candidiasis is very rarely new can cause congenital candidiasis in premature babies (27). Not classified as an infection transmitted by 1; but have frequent sex` It is common in adults (11). Treatment of asymptomatic candidal vulvovaginitis in pregnancy is premature birth` There is tentative evidence that it may reduce the risk of stroke (22).

Diagnosis is made by examining a sample of vaginal mucus. Symptoms are sexual` Infections transmitted by 1, chlamydia and gonorrhea , testing may be recommended (11).

Pregnancy and Vaginal Candidiasis

Studies show that showed that 20-30 percent of women get vaginal candidiasis during pregnancy. Estrogen hormone growth fungus created a favorable environment for colonization of vulvovaginal candidiasis in the vagina, because it binds to protein factor H on the surface of cells that cause vulvovaginal candidiasis will be. But the disease has no effect on the developing fetus does not show. Therefore, treatment is carried out only if candidiasis bothers you. Fungus during pregnancy or breastfeeding Several creams or vaginal suppositories are recommended to treat infections. Over-the-counter medications such as Miconazole, Clotrimazole, and Terconazole treat yeast infections , has been proven to safely and effectively eliminate infection. They usually last three to seven days` is used. It is important to finish the entire course of medication to prevent re-infection. Studies show that showed that these drugs are safe for use during pregnancy (23).

Sex hormones and vaginal candidiasis

Have high estrogen levels The probability of infection with vaginal candidiasis increases. This is due to the high concentration of estrogen in the environment fungal infection white blood cells cannot recognize them and the fungus multiply without any resistance.




Figure 1: Effect of estrogen on *Candida albicans*

- Estrogen increases virulence of *Candida albicans*, reduces phagocytosis
- Estrogen-adapted *C. albicans* o`z binds more factor H on the cell surface

- Binding of estrogen to Gpd2 of *C. albicans*` a family that is immune evasion
- Overexpression of GPD2` growth increases the virulence of *C. albicans* (21).

Diabetes and vaginal candidiasis

In 2018, the data of more than 300,000 people The research that includes showed that people with type 1 or type 2 diabetes have a higher risk of infection, including vaginal candidiasis, than people without diabetes.

Fungus feeds on sugar. In people with diabetes, blood sugar levels can be very high if they are not well controlled`can be strained`growth fungus`of co`can lead to an increase (16).In addition, he has diabetes`conditional infections etc. in sick patients`there is a high risk of organ damage, which impairs their ability to fight pathogens(24). Some more in the body of diabetics`changes, for example: decrease in intestinal secretion, nutritional reasons, gastrointestinal microflora changes, immunodeficiency and coexisting diseases, constant use of antibiotics or other drugs, changes in liver function, lack of necessary nutrients Candida creates a suitable environment for growth (17).

101 diabetic patients of De Leon et found that the frequency of Candida colonization was three times higher in type 1 diabetics compared to type 2 patients. In the same study, the most The most common colonies were Candida albicans, which were detected in 56% of type 1 diabetic patients, while Candida glabrata, in contrast, was found to have a prevalence of 54% of colonies in type 2 diabetic patients.(4).

Table 1

People with diabetes`in healthy patients with VVK` studies comparing lame people

<u>The author</u>	<u>Year</u>	<u>O`studied population groups</u>	<u>Experimental results</u>
Gunther et al	2014	717 women were screened for Candida species`woven; Diabetes group numbers (n); n= 48, Control group n=669	Type 2 diabetes patients in Brazil are healthy significantly more than the lom group more colonization, symptoms and repeated infections they showed 18.8% of diabetics have vaginal Candida species showed, healthy and only 11.8% in disabled people (10).
Goswami et al	<u>2000</u>	166; n=78 Diabetics, n=88 diabetics those who are not.	People with diabetes`46% (36/78) of patients with diabetes had Candida species and only 23% (21/88) of healthy`Candida species in humans`showed(1).
Kendirci and others	2004	57; n=35 type 1 diabetes, n=22 healthy subjects	To`52.5% (32/61) of type 1 diabetes samples showed the presence of Candida species`showed, healthy`only 18.2% (5/22) of the subjects acknowledged its existence`showed (15).
<u>Donders et al</u>	<u>2002</u>	94 women; n=62 women with ≥ three episodes of candidiasis and positive microscopic Candida findings, n=32 for healthy Candida negative controls.	Patient with chronic vulvovaginal candidiasis`36% of patients with diabetes have at least one glucose value above 95%`However, only 12% of the control study population had this finding. HbA1c levels were found to be 25% higher in the recurrent vulvovaginal candidiasis group compared to the control group(5).

AIDS and vaginal candidiasis

Frequency of asymptomatic vaginal colonization o`When studied, it was found that different population groups differ. Candida colonization in HIV negative pregnancy`occurs in 10±20% of women without In

pregnant women, this frequency is up to 15±30%`is narrowed and significantly reduced in post-menopausal women. Women in STD clinics have a higher frequency of asymptomatic colonization. Be pregnant`and be pregnant in non-HIV-positive women conducted a number of studies, microscopy and culture including z, found that vaginal colonization increases significantly in HIV-infected women showed (14,28,30). Also, based on the data, we can say that Candida albicans species prevails in causing vaginal candidiasis even in women with HIV (12).

Vaginal microflora and vaginal candidiasis

Fungus in the vagina In addition to external factors, the vaginal microflora is also of great importance for the diseases. Especially dominant in the vaginal microflora Lactobacillus types. Studies have shown that the dominance of Lactobacillus crispatus and Lactobacillus iners in the vagina has an antagonistic effect on the development of Candida albicans.

In the experiment, 266 women of reproductive age were taken and their vaginal microflora was analyzed:

- L.crispatus dominant in 54 (20%)
- L. iners dominant in 106 people (38 %)
- Other species of Lactobacillus dominate in 106 (42%)

The percentage of chronic vulvovaginal candidiasis was almost 2 times higher in women with other types of Lactobacillus dominating, occupying 42% of it (18).

Treatment

Patients with vaginal infection are treated with saline cream, suppositories and tablets. Medicines usually range from 1-7 days depending on the severity of the disease`is accepted in Treatments are mainly taken at night. The reason for this is keeping creams and suppositories on the affected area longer. Below is the duration of 1 course of treatment of some drugs shown.

Table 2

Medicine	How long to use
Clotrimazole 1% (cream)	7-14 days
Clotrimazole 2% (Cream)	3 days
Miconazole 2% (cream)	7 days
Miconazole 4% (cream)	3 days
Miconazole 100 mg (vaginal suppository)	7 days
Miconazole 200 mg (vaginal suppository)	3 days
Miconazole 400 mg (vaginal suppository)	1 day

Besides that drugs such as butoconazole, clotrimazole, miconazole, nystatin, tioconazole, terconazole are also prescribed (11).

Is it possible to get a vaccine against vaginal candidiasis?

Determination of the immune response in the vagina is the basis for the discovery of vaccination or immunotherapy methods against vaginal candidiasis`ldi Important immunodominant antigen and virulence factors of C.albicans affecting mucosal infections`lgan aspartyl-proteinase (Sap2) with viromas`and received the PEV7 vaccine. In a mouse model and by Pevion in women o`results of a clinical trial suggest that intravaginal PEV7 vaccine is effective for the treatment of recurrent

vulvovaginal candidiasis. that it has a stimulating therapeutic potential showed This is a method of protection against Candida at the mucosal level l opens(8).

Prevention

You often the usual way by changing the vaginal fungus You can prevent infection. This o`changes may include:

- ✓ Use the shower less often - it weakens the vaginal microflora
- ✓ Fragrance`Do not use tampons
- ✓ Use cotton clothes
- ✓ If you have diabetes If so, try to keep your blood sugar in a normal range
- ✓ Always use antifungal agents when taking antibiotics accept
- ✓ If you are pregnant or infected with HIV`If necessary, consult your doctor for treatment (11).

Summary: Vulvovaginal candidiasis is still worldwide`threat to women's health`is standing and numerically it's getting worse. Especially, patients with diabetes and acquired immune deficiency syndrome are more than that`are suffering more. In addition, pregnant women and those with low immunity women are also suffering from this disease. Although the treatment measures are simple, the course of treatment is completed. Incontinence causes vaginal candidosis into a chronic for puts, therefore, be more attentive to treatment we are required. Vaginal candidiasis vaccine has been tested in recent experiments is being made. We hope that this test will be successful. But if everyone tries to prevent the disease, healthy`If you follow a healthy lifestyle, you can avoid vaginal candidiasis and other diseases.

References:

1. Alexander Muacevic and John R Adler`Lubna Mohammed,`Gaurav Jha,`Iana Malasevskaia et al. The Interplay Between Sugar and Yeast Infections: Do Diabetics Have a Greater Predisposition to Develop Oral and Vulvovaginal Candidiasis?`Monitoring Editor:/ 2021 Feb 18. doi:10.7759/cureus.13407
2. Aliyev Sh.R, Muhamedov MI et al. Guide to laboratory training in microbiology 2013.
3. Burns DN, Tuomala R, Chang BH, et al. Vaginal colonization or infection with *Candida albicans* in human immunodeficiency virus-infected women during pregnancy and during the postpartum period. Women and Infants Transmission Study Group. Clin Infect Dis 1997;24:201±1011
4. de Leon EM, Jacober SJ, Sobel JD, Foxman B. Prevalence and risk factors for vaginal *Candida* colonization in women with type 1 and type 2 diabetes. BMC Infect Dis. Jan 30.2002 doi:10.1186/1471-2334-2-1
5. Donders GG, Prenen H, Verbeke G, Reybrouck R. Am J. Impaired tolerance for glucose in women with recurrent vaginal candidiasis. Obstet Gynecol. 2002 PMID: 12388993 DOI:10.1067/mob.2002.126285
6. Duerr A, Sierra MF, Feldman J, Clarke LM, Ehrlich I, DeHovitz J. Immune compromise and prevalence of *Candida* vulvovaginitis in human immunodeficiency virus-infected women. Obstet Gynecol 1997;90:252±69
7. Edwards Jr, JE; Schwartz, MM; Schmidt, CS et al. (2018-06-01). "A Fungal Immunotherapeutic Vaccine (NDV-3A) for Treatment of Recurrent Vulvovaginal Candidiasis-A Phase 2 Randomized, Double-Blind, Placebo-Controlled Trial". Clinical Infectious Diseases. PubMed.gov. 66 (12): 1928–1936. doi:10.1093/cid/ciy185. PMC 5982716. PMID 29697768.)
8. Flavia de Bernardis, Sofia Graziani, Flavio Trelli. *Candida* vaginitis: virulence, host response and vaccine prospects. International society for human and animal mycology. Medical mycology 2018, 56, S26-S31, doi:10.10/93mmy/myx139)
9. Goswami R, Dadhwal V, Tejaswi S, et al. Species-specific prevalence of vaginal candidiasis among patients with diabetes mellitus and its relation to their glycemic status. J Infect. 2000 Sep;41(2):162-6. doi: 10.1053/jinf.2000.0723

10. Gunther LS, Martins HP, Gimenez F, de Abreu ALP, Consolaro MEL, Estivalet TI Prevalence of *Candida albicans* and non-*albicans* isolates from vaginal secretions: comparative evaluation of colonization, vaginal candidiasis and recurrent vaginal candidiasis in diabetic and non-diabetic women. . Sao Paulo Med J.2014;132(2):116-20.doi:10.1590/15163180.2014.1322640
11. Hubertine ME Willems, Salman S. Ahmed, Junyan Liu, Zhenbo Huo, Brian M. Peters Vulvovaginal Candidiasis: A Current Understanding and Burning Questions 2020 Feb 25. doi:10.3390/jof6010027
12. Jack D Sobel MD Chief Vulvovaginal candidiasis: a comparison of HIV-positive and -negative women , Division of Infectious Diseases, Department of Internal Medicine, Wayne State University School of Medicine, Detroit, MI, International Journal of STD & AIDS 2002;13: 358
13. Jane Carpenter, Ethel Burns, Lesley Smith Journal of midwifery and women's health Nov. 2021 doi:10.1111/jmwh.133326)
14. Kapiga SH, Lyamuya EF, Lwihula GK, Hunter DJ. The incidence of HIV infection among women using family planning methods in Dares Salaam, Tanzania. AIDS 1998;12:75±8410
15. Kendirci M, Koç AN, Kurtoglu S, Keskin M, Kuyucu T. J Pediatr Endocrinol Metab. Vulvovaginal candidiasis in children and adolescents with type 1 diabetes mellitus. 2004; doi: 10.1515/jpem.2004.17.11.1545
16. M. Carey Julia A. Critchley; Stephen DeWilde; Tess Harris; Fay J. Hosking; Derek G. Cook Corresponding author: Iain M. Carey, i.carey@sgul.ac.uk. Diabetes Care 2018;41(3):513–521 <https://doi.org/10.2337/dc17-2131>)
17. Martins N, Ferreira IC, Barros L, Silva S, Henriques M. Candidiasis: predisposing factors, prevention, diagnosis and alternative treatment. Mycopathology. 2014;177:223–240
18. Mr. Brett A. TORTELLI, BA, Warren G. LEWIS, Ph.D., Jennifer E. ALLSWORTH, Ph.D., Mr. Nadum MEMBER-MENEH, BS, Ms. Lynne R. FOSTER, BS, Hilary E. RENO, MD Ph.D., Jeffrey F. PEIPERT, MD, Ph.D., Justin C. FAY, Ph.D. oath Amanda L. LEWIS, Ph.D. Associations between the vaginal microbiome and *Candida* colonization in women of reproductive age 2019 Oct 22. doi:10.1016/j.ajog.2019.10.008
19. Nicodemus, Eva; Tamasi, Béla; Mihalik, Noémi; Ostorházi, Eszter (1 January 2015). "Yeast species in vulvovaginitis candidosa". Orvosi Hetilap (in Hungarian). 156 (1): 28–31. doi:10.1556/OH.2015.30081. PMID 25544052.
20. Pappas, PG; Kauffman, CA; Andes, DR; Clancy, CJ; Marr, KA; Ostrosky-Zeichner, L; Reboli, AC; Schuster, MG; Vazquez, JA; Walsh, TJ; Zaoutis, TE; Sobel, JD (16 December 2015). "Clinical Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious Diseases Society of America". Clinical Infectious Diseases. 62(4):e1-50. doi:10.1093/cid/civ933. PMC 4725385. PMID 26679628)
21. Pizga Kumwenda, Fabien Cottier, Alexandra C. Hendry, Davey Kneafsey, Ben Keevan, 1 Hannah Gallagher, 1 Hung-Ji Tsai, 1 and Rebecca A. Hall 1. 1Institute of Microbiology and Infection, School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK 2Kent Fungal Group, Division of Natural Sciences, School of Biosciences, University of Kent, Canterbury CT2 7NJ, UK 3Lead contact *Correspondence: rahall@kent.ac.uk <https://doi.org/10.1016/j.celrep.2021.110183>
22. Roberts, CL; Algert, CS; Rickard, KL; Morris, JM (21 March 2015). "Treatment of vaginal candidiasis for the prevention of preterm birth: a systematic review and meta-analysis". Systematic Reviews. 4: 31. doi:10.1186/s13643-015-0018-2. PMC 4373465. PMID 25874659.]
23. Robyn Horsager, Boehrer MD Obstetrics and Gynecology ecology/DOI:10.1097/00006250-199406000-0001120. September 20, 2022)
24. Rodrigues CF, Rodrigues ME, Henriques M. J Candida sp. infections in patients with diabetes mellitus. Clin Med. 2019; 8:76.
25. Schuman P, Sobel JD, Ohmit SE, et al. Mucosal candidal colonization and candidiasis in women with or at risk for human immunodeficiency virus infection. HIV Epidemiology Research Study (HERS) Group. Clin Infect Dis 1998;27:1161±78

26. Seyedmojtaba Seyedmousavi, Sandra de MG Bosco, Sybren de Hoog, Frank Ebel, Daniel Elad, Renata R Gomez, Ilse D Jacobsen, Henrik E Jensen, Anne Martel, Bernard Mignon Medical Mycology, Volume 56, Issue suppl_1, April 2018, Pages S26–S31, <https://doi.org/10.1093/mmy/myx139>
27. Skoczylas, MM; Walat, A; Kordek, A; Loniewska, B; Rudnicki, J; Maleszka, R; Torbé, A (2014). "Congenital candidiasis as a subject of research in medicine and human ecology". *Annals of Parasitology*. 60 (3): 179–89. PMID 25281815.)
28. Sobel JD, Ohmit SE, Schuman P, et al. The evolution of *Candida* spp. and uconazole susceptibility among oral and *Infect Dis* 2001;183:286±9312
29. Sobel, JD (9 June 2007). "Vulvovaginal candidiasis". *Lancet*. 369 (9577): 1961–71. doi:10.1016/S0140-6736(07)60917-9. PMID 17560449. S2CID 33894309.]
30. Spinillo A, Michelone G, Cavanna C, Colonna L, Capuzzo E, Nicola S. Clinical and microbiological characteristics of symptomatic vulvovaginal candidiasis in HIV-seropositive women. *Genitourin Med* 1994;70:268±72
31. Workowski KA, Berman SM (August 2006). "Sexually transmitted diseases treatment guidelines, 2006". *MMWR Recomm Rep*. 55 (RR-11): 1–94. PMID 16888612. Archived from the original on 2014-10-20.
32. Molchanov Oleg Leonidovich. Intimate hygiene during pregnancy "EGIS-RUS" 2021/ PN015678/01
33. Pestrikova T. Yu., Yurasova E.A. Kotelnikova A.V. Vulvovaginal candidiasis: a modern perspective and problem. *RMJ (Russian medical journal)*. 2018. LP 004092-230117